

DNA purification system

- Production of cleared lysate
 - Pellet 1-10ml of overnight culture for 5 minutes
 - Thoroughly resuspend pellet with 250µl of Cell Resuspension Solution
 - Add 250µl of Cell Lysis Solution to each sample; invert 4 times to mix
 - Add 10µl of Alkaline Protease Solution; invert 4 times to mix. Incubate 5 minutes at room temperature
 - Add 350µl of Neutralization Solution; invert 4 times to mix
 - Centrifuge at top speed for 10 minutes at room temperature
- Binding of plasmid DNA
 - Insert Spin Column into Collection Tube
 - Decant cleared lysate into Spin Column
 - Centrifuge at top speed for 1 minute at room temperature. Discard flowthrough, and reinsert Column into Collection Tube.
- Washing
 - Add 750µl of Wash Solution (ethanol added). Centrifuge at top speed for 1 minute. Discard flowthrough and reinsert column into Collection Tube
 - Repeat step before with 250µl of Wash solution
 - Centrifuge at top speed for 2 minutes at room temperature
- Elution
 - Transfer Spin Column to a sterile 1.5ml microcentrifuge tube, being careful not to transfer any of the Column Wash Solution with the Spin Column. If the Spin Column has Column Wash Solution associated with it, centrifuge again for 1 minute at top speed, then transfer the Spin Column to a new, sterile 1.5ml microcentrifuge tube
 - Add 100µl of Nuclease-Free Water to the Spin Column. Centrifuge at top speed for 1 minute at room temperature
 - Discard column, and store DNA at -20°C or below

